

The Mass Gap of the Nonlinear σ Model through the Finite Temperature Effective Action

David Sénéchal

*Centre de Recherche en Physique du Solide et Département de Physique,
Université de Sherbrooke, Sherbrooke, Québec, Canada J1K 2R1.*

(November 1992)

Abstract

The $O(3)$ nonlinear σ model is studied in the disordered phase, using the techniques of the effective action and finite temperature field theory. The nonlinear constraint is implemented through a Lagrange multiplier. The finite temperature effective potential for this multiplier is calculated at one loop. The existence of a nontrivial minimum for this potential is the signal of a disordered phase in which the lowest excited state is a massive triplet. The mass gap is easily calculated as a function of temperature in dimensions 1, 2 and 3. In dimension 1, this gap is known as the Haldane gap, and its temperature dependence is compared with experimental results.

75.10.Jm, 11.10.Lm

The $O(3)$ nonlinear σ model (NL σ) offers a good description of quantum antiferromagnets at long wavelengths, as suggested by renormalization group analyses¹. The mapping between the nonlinear σ model and the Heisenberg model was first established by Haldane², who also argued that a topological Hopf term should be present in the case of half-integer spins in one dimension³ (this distinction between half-integer and integer spins disappears in two dimensions). For integer spins, in which case the Hopf term is absent, Haldane argued that the lowest excitations should exhibit a mass gap. Accordingly, these excitations (the magnons) are no longer Goldstone modes and the rotation symmetry of the spins is no longer spontaneously broken: the symmetry has been dynamically restored by quantum fluctuations. This phenomenon has found an experimental realization in quasi one-dimensional antiferromagnets in their disordered phase⁴.

In this paper we calculate the temperature dependence of this mass gap in dimensions 1 to 3 within the framework of the finite temperature effective action of the nonlinear σ model. The use of this technique is facilitated by the Lorentz invariance of the NL σ model, which ensures that the magnon speed is not affected by quantum corrections.

The $O(3)$ nonlinear σ model describes the dynamics of a unit vector $\vec{\varphi}$ representing the local direction of the staggered magnetization. The Euclidian action obtained from the Heisenberg model is

$$S = \frac{v}{2g} \int d\tau d^d x \left\{ \frac{1}{v^2} (\partial_\tau \vec{\varphi})^2 + (\partial_i \vec{\varphi})^2 \right\} \quad (1)$$

wherein v is the spin-wave velocity and g a coupling constant. the constraint $\vec{\varphi}^2 = 1$ is understood. If we choose a set of units in which the speed v is unity, implement the constraint $\vec{\varphi}^2 = 1$ by a Lagrange multiplier σ , and rescale the field $\vec{\varphi}$ by $\sqrt{g}$ in order to bring its kinetic term into the standard form, we obtain the equivalent action

$$S = \int d\tau d^d x \left\{ \frac{1}{2} \partial_\mu \vec{\varphi} \partial_\mu \vec{\varphi} - \frac{1}{2} \sigma (\vec{\varphi}^2 - 1/g) \right\} \quad (2)$$

wherein the index μ goes from 0 to d . The spin-wave velocity v may always be restored later using dimensional analysis.

The action (2) is symmetric with respect to global rotations of the field $\vec{\varphi}$, which is the relevant order parameter. The degenerate classical ground states are configurations in which the field $\vec{\varphi}$ is uniform. Such ground states break the full symmetry group $O(3)$ down to $O(2)$. Goldstone's theorem would then imply the existence of two massless modes, corresponding to the two magnon polarizations. However, the quantum ground state, to which Goldstone's theorem refers, does not break the $O(3)$ symmetry if we are in a disordered phase. The symmetry of the quantum ground state may be determined by calculating the one-loop effective potential $V(\sigma)$ for the field σ , following the discussion by Coleman⁵. If this potential has a global minimum at some nonzero value σ_0 of the field σ , then we must expand the field σ around its ground state value σ_0 . The latter then gives rise to a mass term $\frac{1}{2}\sigma_0\vec{\varphi}^2$ in the action (2). As a consequence, the lowest lying excitations are no longer a pair of massless Goldstone modes (the infrared instability has been removed) but a massive triplet: the elementary excitations of the three-component field $\vec{\varphi}$ with mass gap $\Delta = \sqrt{\sigma_0}$.

The one-loop effective potential $V(\sigma)$ is obtained by summing an infinite sequence of Feynman graphs, each with n insertions of σ and n internal $\vec{\varphi}$ lines. This amounts to evaluating the partition function with σ constant, which is done simply by Gaussian integration. The result of this standard calculation is

$$V(\sigma) = \frac{\sigma}{2g} - 3 \int \frac{dp_0}{(2\pi)} \int \frac{d\vec{p}}{(2\pi)^d} \log \left\{ 1 + \frac{\sigma}{p_0^2 + \vec{p}^2} \right\} \quad (3)$$

where we have included the tree-level potential $\sigma/2g$ (we use the convention $\hbar = 1$ throughout). In order to find out whether $V(\sigma)$ has a minimum, we consider instead the derivative

$$V'(\sigma) = \frac{1}{2g} - 3 \int \frac{dp_0}{(2\pi)} \int \frac{d\vec{p}}{(2\pi)^d} \frac{1}{p_0^2 + \vec{p}^2 + \sigma} \quad (4)$$

The extension of this formula to a finite temperature T is obtained by replacing the integration over p_0 by a sum over Matsubara frequencies $\omega_n = 2\pi nT$:

$$\int \frac{dp_0}{2\pi} f(p_0) \rightarrow T \sum_n f(\omega_n) \quad (5)$$

Using the standard result ($\beta = 1/T$)

$$\sum_n \frac{1}{\omega_n^2 + z^2} = \frac{\beta}{2z} \coth(\beta z/2) \quad (6)$$

we may write

$$2V'(\sigma) = \frac{1}{g} - 3 \int \frac{d\vec{p}}{(2\pi)^d} \frac{1}{\sqrt{\vec{p}^2 + \sigma}} \coth\left(\frac{1}{2}\beta\sqrt{\vec{p}^2 + \sigma}\right) \quad (7)$$

At zero temperature, the above reduces to

$$2V'(\sigma) = \frac{1}{g} - 3K_d \int_0^\Lambda \frac{p^{d-1} dp}{\sqrt{p^2 + \sigma}} \quad (8)$$

where $K_1^{-1} = 2\pi$ and $K_d^{-1} = 2^{d-1}\pi^{d/2}\Gamma(d/2)$ if $d > 1$. An ultraviolet cutoff Λ has been introduced. We point out that the NL σ model is a renormalizable field theory only in $d = 1$; this means that our treatment for $d > 1$ will only be valid in the limit of large distances. Without taking Λ to infinity, we will consistently neglect terms of order Λ^{-1} or more. This being understood, finite temperatures will not introduce any new dependence on Λ : Indeed, the difference $V_\delta \equiv V(\sigma, T) - V(\sigma, 0)$ is given by

$$2V'_\delta(\sigma) = -3K_d \int_0^\Lambda \frac{p^{d-1} dp}{\sqrt{p^2 + \sigma}} \left\{ \coth\left(\frac{1}{2}\beta\sqrt{p^2 + \sigma}\right) - 1 \right\} \quad (9)$$

Since the function $\coth x - 1$ decreases exponentially* as $x \rightarrow \infty$, the above integral is UV finite and the cut-off may be removed. The dependence on Λ is contained solely in $V(\sigma, 0)$. A change of integration variables yields

$$2V'_\delta(\sigma) = -T^{d-1} f_d(y) \quad (10)$$

where $y = \sqrt{\sigma}/T$ and

$$f_d(y) = 3y^{d-1} K_d \int_1^\infty dz \frac{\coth(\frac{1}{2}yz) - 1}{(z^2 - 1)^{1-d/2}} \quad (11)$$

We find that

$$f_1(y) = \frac{3}{\pi} \sum_{n=1}^\infty K_0(ny) \quad (12a)$$

$$f_2(y) = \frac{3}{\pi} \left\{ \frac{1}{2}y - \log(2\sinh(y/2)) \right\} \quad (12b)$$

$$f_3(y) = \frac{3y}{\pi^2} \sum_{n=1}^\infty \frac{1}{n} K_1(ny) \quad (12c)$$

(K_0 and K_1 are modified Bessel functions.)

Let us now look in more detail at specific cases, starting with $d = 1$. This has been done explicitly by Coleman⁵ for the CP^1 model, which is equivalent to the $O(3)$ nonlinear σ model. From Eq. (8) we obtain, at zero temperature,**

$$2V'(\sigma) = \frac{1}{g} + \frac{3}{2\pi} \log(\sigma/\Lambda^2) \quad (d = 1) \quad (13)$$

We see that the equation $V'(\sigma) = 0$ always has a solution, at the point $\sigma_0 = \Lambda^2 \exp -(2\pi/3g)$, which implies a zero temperature Haldane gap $\Delta_0 = \Lambda \exp -\pi/3g$.

From Eqs. (10) and (12), the finite temperature result is then

$$2V'(\sigma) = \frac{1}{g} + \frac{3}{2\pi} \log(\sigma/\Lambda^2) - \frac{3}{\pi} \sum_{n=1}^{\infty} K_0(n\sqrt{\sigma}/T) \quad (14)$$

The finite temperature Haldane gap $\Delta = \sqrt{\sigma_0}$ is therefore obtained through the solution of the equation

$$\frac{1}{2} \log(\delta) = \sum_{n=1}^{\infty} K_0(n\delta/t) \quad (15)$$

where we have defined the reduced gap $\delta = \Delta/\Delta_0$ and the reduced temperature $t = T/\Delta_0$. This gap equation may be solved numerically. The solution is shown on Fig.1. The gap is seen to maintain its $T = 0$ value until about $T \sim 0.2\Delta_0$, after which it takes off linearly. We naturally expect the gap to grow with temperature since Δ , being the inverse of the correlation length, is in some sense a measure of disorder. This temperature dependence seems confirmed by experiments⁶.

The dependence of Δ on T is not qualitatively affected by the introduction of anisotropy in the σ model. Easy-axis anisotropy is introduced via a mass term $\frac{1}{2}M^2\varphi_z^2$ in the action (2), while easy-plane anisotropy is rendered by a mass term $\frac{1}{2}M^2(\varphi_x^2 + \varphi_y^2)$. The above calculation of the Haldane gap is easily adapted to the anisotropic case by shifting σ by α for one (two) out of three components of $\vec{\varphi}$ in the easy-axis (easy-plane) case. The gap equation (15) is thus modified into

$$\frac{1}{4} \log \left\{ \left(\frac{\mu^2 + \delta^2}{\mu^2 + 1} \right) \delta^4 \right\} = \sum_{n=1}^{\infty} K_0(n\sqrt{\delta^2 + \mu^2}/t) + 2 \sum_{n=1}^{\infty} K_0(n\delta/t) \quad (16)$$

where we have defined the reduced anisotropy $\mu = M/\Delta_0$ and considered the easy-axis case. Even if the value of Δ_0 is affected by the anisotropy, everything else being fixed, we have checked that the t dependence of the reduced gap δ is not changed appreciably by the introduction of an anisotropy even as large as $\mu = 1$. However, an easy-axis anisotropy splits the Haldane triplet into a doublet with gap $\Delta_1 = \Delta$ and a singlet with gap $\Delta_2 = \sqrt{\Delta^2 + M^2}$. The weighted average of the two gaps, normalized to its zero temperature value, is shown on the lower curve of Fig.1 for $\mu = 1.9$. This value of μ , appropriate for NENP, is obtained from the observed values of Δ_1 (14K) and Δ_2 (30K)⁷.

We next consider the two-dimensional case. At $T = 0$ this was studied by Deser and Redlich⁸ for the CP^1 model. The result of the integration (8) is

$$2V'(\sigma) = \frac{1}{g} - \frac{3\Lambda}{2\pi} + \frac{3\sqrt{\sigma}}{2\pi} \quad (d = 2) \quad (17)$$

The interesting feature here is the appearance of a critical coupling $g_c = 2\pi/\Lambda$ below which the gap equation $V'(\sigma) = 0$ has no solution. This is the signal of a phase transition in coupling space: below g_c , the $O(3)$ symmetry is spontaneously broken and massless Goldstone modes appear. At finite temperatures we have, following Eq. (12):

$$2V'(\sigma) = \frac{1}{g} - \frac{1}{g_c} + \frac{3T}{\pi} \log \left[2 \sinh(\sqrt{\sigma}/2T) \right] \quad (18)$$

The gap equation has now a solution for all nonzero values of T , indicating that the ordered phase does not survive at finite temperatures, as is already known from RG calculations¹.

Let us define the difference $\eta = 1/g_c - 1/g$. The solution of the gap equation is then

$$\Delta = 2T \sinh^{-1} \left[\frac{1}{2} \exp(\pi\eta/3T) \right] \quad (19)$$

If $\eta > 0$ (disordered phase) the gap tends towards the constant $2\pi\eta/3$ as $T \rightarrow 0$, i.e., $\eta = 3\Delta_0/2\pi$. If, on the contrary, $\eta < 0$ (ordered phase) the gap behaves like $\Delta \approx T \exp -\pi|\eta|/3T$ as $T \rightarrow 0$, i.e., it vanishes faster than any power law. At the critical value $\eta = 0$ the gap vanishes linearly as T is lowered:

$$\Delta = 2T \sinh^{-1}(\frac{1}{2}) \approx 0.962 T \quad (20)$$

These results are illustrated in Fig.2. Similar results were obtained within the RG analysis of Ref.[1] in terms of the correlation length $\xi = 1/\Delta$. Experimental results are available in the ordered phase ($\eta < 0$) and compare well with the theory (see Ref.[1]). To the author's knowledge, the disordered phase ($\eta > 0$) has not been observed yet. Although it is accepted that this phase does not exist at $T = 0$ for the Heisenberg model with spin $s \geq 1$, nothing certain can be said about the spin- $\frac{1}{2}$ case.

Let us point out that the cross-over region, i.e., the temperature range over which thermal fluctuations overtake quantum fluctuations, is characterized by the condition $\Delta \sim T$ which, on the phase diagram (g, T) , is met on the curve $T \sim 25.3\eta$, according to Eq. (19). When T goes beyond this value the states over the gap are thermally excited or, stated differently, the correlation length ξ exceeds the extent β of the Euclidian time direction; the fluctuations then happen mostly in the spatial directions.

Let us finally discuss the three-dimensional case, first at $T = 0$. The integration (8) yields

$$2V'(\sigma) = \frac{1}{g} - \frac{3\Lambda^2}{4\pi^2} + \frac{3\sigma}{8\pi^2} \left\{ \log(2\Lambda^2/\sigma) - 1 \right\} \quad (21)$$

Again, we have dropped terms of order $1/\Lambda$. It is understood that σ is much smaller than the cutoff Λ , so that $V'(\sigma)$ is increasing monotonically with σ . Like in dimension 2, there is a critical coupling $g_c = 4\pi^2/3\Lambda^2$ below which the symmetry is spontaneously broken at $T = 0$. The difference with $d = 2$ is that this ordered phase survives at finite temperature. Indeed, following Eqs. (12) and (21), the gap equation is

$$0 = \frac{1}{g} - \frac{1}{g_c} + \frac{3\Delta^2}{8\pi^2} \left\{ \log(2\Lambda^2/\Delta^2) - 1 \right\} - T^2 f_3(\Delta/T) \quad (22)$$

The function $f_3(y)$ is monotonically decreasing, rather like an exponential, starting with $f_3(0) \approx 0.5$. If the above equation has no solution at $T = 0$ ($g < g_c$), then an increase in temperature will bring a solution at some point, since the r.h.s. can only decrease as T increases. The transition to the ordered phase occurs when $\Delta = 0$; the relation between g and T that follows is

$$\frac{1}{g} - \frac{1}{g_c} = T^2 f_3(0) \quad (23)$$

We see that no matter how small g may be below g_c , a critical temperature exists beyond which we recover a disordered phase. However, this result is not to be trusted at arbitrary low g , since we tacitly assumed that $T \ll \Lambda$.

If we place ourselves above g_c , denote by Δ_0 the zero temperature gap and define reduced variables $\delta = \Delta/\Delta_0$, $t = T/\Delta_0$ and $A = \log(\Lambda/\Delta_0) + \frac{1}{2}(\log 2 - 1)$, the gap equation becomes

$$A(1 - \delta^2) - \delta^2 \log \delta + \frac{4\pi}{3} t^2 f_3(\delta/t) = 0 \quad (24)$$

Its solution for δ as a function of t may be found numerically, and is shown on Fig. 3 for two values of A . Note that we cannot rid ourselves of this extra parameter, but its logarithmic dependence on the cut-off causes only a mild non universality. At the critical value $g = g_c$, we find that the gap vanishes linearly with T , like in dimension 2, albeit with mild corrections.

In conclusion, calculating the one-loop effective potential for the Lagrange multiplier σ yields the Haldane gap as a function of temperature in a rather direct way. The solution is compatible with experimental results. The extension to two and three dimensions is straightforward although, to the author's knowledge, experimental results in the quantum disordered regime are then lacking. Since no rigorous argument has yet been found that prevents the existence of a quantum disordered phase in the two-dimensional spin- $\frac{1}{2}$ Heisenberg model, the question remains open.

ACKNOWLEDGMENTS

Stimulating conversations with C. Bourbonnais, W.J.L. Buyers, M. Plumer, Y. Trudeau and especially A.-M.S. Tremblay are gratefully acknowledged. This work is supported in part by le Fonds pour la Formation de Chercheurs et l'Aide à la Recherche du Gouvernement du Québec (F.C.A.R.).

REFERENCES

- * The expansion $\coth x - 1 = 2 \sum_{n=1}^{\infty} \exp -2nx$ has proven useful.
- ** The logarithmic cut-off dependence could be removed by defining a renormalized coupling : $1/g_r \equiv 2V'(\mu^2)$, where μ is a subtraction point. The effective potential would then be well-behaved as $\Lambda \rightarrow \infty$ and $g \rightarrow 0$, while g_r stays fixed. However, since we do not intend to send the cutoff to ∞ , this procedure would be of little use to us.
- ¹ S. Chakravarty, B. I. Halperin and D. R. Nelson, Phys. Rev. **B39**, 2344 (1989).
- ² F. D. M. Haldane, Phys. Lett. **A93**, 464 (1983).
- ³ F. D. M. Haldane, Phys. Rev. Lett. **50**, 1153 (1983).
- ⁴ W. J. L. Buyers et al., Phys. Rev. Lett. **56**, 371 (1986).
- ⁵ S. Coleman, *Aspects of Symmetry: selected Erice Lectures*, Cambridge University Press (1985), chapter 8.
- ⁶ W. J. L. Buyers, Z. Tun, A. Harrison, J. A. Rayne and R. M. Nicklow, Physica **B180-181**, 222 (1992).
- ⁷ J.-P. Renard et al., J. Appl. Phys. **63**, 3538 (1988).
- ⁸ S. Deser and A. N. Redlich, Phys. Rev. Lett. **61**, 1541 (1988).

FIGURES

FIG. 1. Magnitude of the reduced Haldane gap $\delta = \Delta/\Delta_0$ as a function of $t = T/\Delta_0$ in $d = 1$ (upper curve). Experimental results from Ref.[6] are also shown for CsNiCl_3 and NENP (error bars are not included). The lower curve is the weighted average of the two gaps in the presence of a $\mu = 1.9$ easy-axis anisotropy, appropriate for NENP.

FIG. 2. Magnitude of the mass gap Δ as a function of temperature for $d = 2$ and three values (indicated) of η .

FIG. 3. Magnitude of the reduced mass gap $\delta = \Delta/\Delta_0$ as a function of $t = T/\Delta_0$ in $d = 3$ for $A = 4$ (higher curve) and $A = 5$ (lower curve).